

My daily medication tracker



Use this medication tracker daily to help you stay on top of your treatment with TRUQAP. Here's how to get started:

- In the top row, write the date below each day
- Write the time you take each AM and PM dose of TRUQAP in the appropriate box



Things to remember

- TRUQAP is taken 4 days in a row, followed by 3 days not taking TRUQAP, each week
- TRUQAP is taken 2 times a day about 12 hours apart
- Take your morning and night doses of TRUQAP at the same time each day
- If you miss a dose:
 - You may still take it within 4 hours from the time you usually take it
 - If it has been more than 4 hours after you usually take your dose, skip that dose and take your next dose at your usual time
 - Do not take 2 doses at once to make up for a missed dose
- If you vomit after taking a dose, do not take another dose. Take your next dose at your usual time
- Do not chew, crush, or split tablets prior to swallowing. Do not take any tablets that are broken, cracked, or that look damaged

IMPORTANT PRODUCT INFORMATION

TRUQAP® (capivasertib) is a prescription oral medicine used in combination with fulvestrant to treat adults who have hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer that is advanced or has spread to other parts of the body (metastatic), with an abnormal "PIK3CA", "AKT1", or "PTEN" gene, and whose disease has progressed on or after endocrine therapy. Do not take TRUQAP if you have had a severe allergic reaction to TRUQAP or are allergic to any of its ingredients. Tell your healthcare provider if you are pregnant or breastfeeding or plan to become pregnant or breastfeed. TRUQAP can harm your unborn baby. TRUQAP may cause severe side effects, including high blood sugar levels (hyperglycemia); diarrhea; or skin reactions. Untreated severe hyperglycemia can lead to a condition called diabetic ketoacidosis that requires treatment in a hospital and can lead to death. The most common side effects include nausea, tiredness, vomiting, mouth sores, and changes in certain blood tests.

Please see full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.

You may [report side effects related to AstraZeneca products](#).

My daily medication tracker



TRUQAP is taken 4 days in a row, followed by 3 days not taking TRUQAP, each week. TRUQAP is taken 2 times a day about 12 hours apart. Take your morning and night doses of TRUQAP at the same time each day. The recommended starting daily dose is 800 mg; two 200 mg tablets twice a day (400 mg in the morning and 400 mg at night). Take TRUQAP exactly as your doctor tells you to, as your doctor may adjust your dose. Do not change your dose or stop taking TRUQAP unless your doctor tells you to.

DATE	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
 AM 00	:	:	:	:	No Dose	No Dose	No Dose
 PM 00	:	:	:	:			

DATE	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
 AM 00	:	:	:	:	No Dose	No Dose	No Dose
 PM 00	:	:	:	:			

DATE	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
 AM 00	:	:	:	:	No Dose	No Dose	No Dose
 PM 00	:	:	:	:			

DATE	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
 AM 00	:	:	:	:	No Dose	No Dose	No Dose
 PM 00	:	:	:	:			



To download your custom monthly dosing calendar, visit <https://www.truqap.com/about/taking-truqap#calendar>



TRUQAP is a registered trademark of the AstraZeneca group of companies.
©2025 AstraZeneca. All rights reserved. US-96077 Last Updated 2/25