

For people with HR+/HER2- metastatic breast cancer with abnormal *PIK3CA*, *AKT1*, and/or *PTEN* genes following progression on or after hormone therapy.

UNFOLDING MORE

OF YOUR STORY

MAY BE POSSIBLE*



Not an actual patient.

The trial showed no significant difference in how long patients lived between treatment groups.

*TRUQAP + fulvestrant delivered an average of 7.3 months without cancer growing or spreading vs 3.1 months with fulvestrant alone—a difference of 4.2 months. Results may vary.

IMPORTANT SAFETY INFORMATION ABOUT TRUQAP® (capivasertib) tablets

Do not take TRUQAP if you have had a severe allergic reaction to TRUQAP or are allergic to any of the ingredients of TRUQAP.

Before you take TRUQAP, tell your healthcare provider about all your medical conditions, including if you:

- have a history of high levels of sugar in your blood or diabetes
- have a history of rash or other skin disorders
- have liver problems
- have any current infection
- are pregnant or plan to become pregnant. TRUQAP can harm your unborn baby

Please see additional Important Safety Information throughout, and full Prescribing Information, including Patient Information for TRUQAP.

 **Truqap®**
capivasertib
160 mg • 200 mg tablets



Glossary

Throughout this brochure, you will see abbreviations for some of the terms that are associated with HR+/HER2- metastatic breast cancer. Here's what they mean:

AKT1=serine/threonine protein kinase 1

CDK4/6=cyclin-dependent kinase 4/6

HER2-=human epidermal growth factor receptor 2 negative

HR+=hormone receptor positive

PIK3CA=phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha

PTEN=phosphatase and tensin homolog

Not an actual patient.

IMPORTANT SAFETY INFORMATION (continued)

Before you take TRUQAP, tell your healthcare provider about all your medical conditions, including if you: (continued)

Females who are able to become pregnant:

- Your healthcare provider will check to see if you are pregnant before you start treatment with TRUQAP
- You should use effective birth control during treatment with TRUQAP and for 1 month after the last dose. Talk to your healthcare provider about birth control methods that may be right for you during this time
- If you become pregnant or think you are pregnant, tell your healthcare provider right away

Males with female partners who are able to become pregnant should use effective birth control during your treatment with TRUQAP and for 4 months after the last dose. If your female partner becomes pregnant, tell your healthcare provider right away.

- are breastfeeding or plan to breastfeed. Do not breastfeed during treatment with TRUQAP. Talk to your healthcare provider about the best way to feed your baby during treatment with TRUQAP

Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.



Biomarker testing & how TRUQAP works



TRUQAP is for people with HR+/HER2- metastatic breast cancer that tests positive for abnormal *PIK3CA*, *AKT1*, and/or *PTEN* genes

When you were diagnosed, your doctor may have tested you for different biomarkers. Biomarkers can show if there are abnormal changes to the genes in your tumor. These changes are called mutations or alterations. Gene mutations or alterations can contribute to the growth and spread of your cancer. Knowing if your cancer has a targetable biomarker can help determine which treatments may be right for you.

TRUQAP works in people whose cancer has biomarker testing that shows abnormal changes in *PIK3CA*, *AKT1*, and/or *PTEN* genes—and whose disease has progressed on or after hormone therapy. If you haven't been tested for biomarkers yet, you can still be tested at any time. Ask your doctor about biomarker testing at your next appointment to see if TRUQAP could be an option for you.

TRUQAP is the first therapy of its kind

When it's time for you to start your next treatment, TRUQAP + fulvestrant may be right for you

In people with HR+/HER2- metastatic breast cancer with abnormal *PIK3CA*, *AKT1*, and/or *PTEN* genes, overactive signaling in cancer cells can make the tumor grow. TRUQAP is an AKT inhibitor that works by disrupting signals from the AKT protein and other proteins along its pathway that can drive cancer growth.

TRUQAP is taken with the hormone therapy fulvestrant. Fulvestrant works by disrupting signals from the hormone estrogen that fuels tumor growth.

These effects have only been demonstrated in the laboratory in cell studies.

TRUQAP + fulvestrant work together to disrupt the signals that drive cancer growth



Find out if TRUQAP could be right for you.
Talk to your doctor about biomarker testing.

Stay on hormone therapy longer
with TRUQAP + fulvestrant



For people with HR+/HER2- metastatic breast cancer with an abnormal *PIK3CA*, *AKT1*, and/or *PTEN* gene following progression on or after hormone therapy,

TRUQAP + fulvestrant keeps cancer from growing or spreading for **more than 2X longer** than fulvestrant alone

TRUQAP + fulvestrant is proven to give people with HR+/HER2- metastatic breast cancer with abnormal *PIK3CA*, *AKT1*, and/or *PTEN* genes more than 2X longer without their cancer growing or spreading compared with the hormone therapy fulvestrant alone. This is called progression-free survival.

**MORE
THAN 2X**

**LONGER WITHOUT
TUMOR GROWTH**

TRUQAP + fulvestrant delivered an average of 7.3 months vs 3.1 months with fulvestrant alone—a difference of 4.2 months.*

Results may vary.

The trial showed no significant difference in how long patients lived between treatment groups.

*Median time from the start of treatment that patients' tumors did not grow or spread.
Median is the middle number in a group of numbers listed from lowest to highest.

Not an actual patient.

IMPORTANT SAFETY INFORMATION (continued)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. TRUQAP may affect the way other medicines work, and other medicines may affect how TRUQAP works. Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

How should I take TRUQAP?

- Take TRUQAP exactly as your healthcare provider tells you
- Do not change your dose or stop taking TRUQAP unless your healthcare provider tells you

Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.

How TRUQAP was studied

TRUQAP was studied in a global clinical trial that included 708 people with HR+/HER2- metastatic breast cancer with or without abnormal *PIK3CA*, *AKT1*, and/or *PTEN* genes whose cancer grew or spread on hormone therapy.

708
People

355
TRUQAP +
fulvestrant

353
Placebo +
fulvestrant

- Pre/peri- and postmenopausal women, and men*
- Average age of 58
- All patients had received prior hormone (aromatase inhibitor) therapy for their breast cancer
- 70% of people in the study had been previously treated with a CDK4/6 inhibitor

FDA approval was based on results in 289 patients with abnormal *PIK3CA*, *AKT1*, and/or *PTEN* genes (155 patients treated with TRUQAP + fulvestrant vs 134 patients treated with placebo + fulvestrant).



TRUQAP is for people with HR+/HER2- metastatic breast cancer that tests positive for abnormal *PIK3CA*, *AKT1*, and/or *PTEN* genes.

*Pre-/perimenopausal women were also given a luteinizing hormone-releasing hormone agonist.

TRUQAP + fulvestrant keeps cancer from growing or spreading for more than 2X longer vs fulvestrant alone

HOW TO TAKE
TRUQAP
+ FULVESTRANT






Not an actual patient.
The trial showed no significant difference in how long patients lived between treatment groups.

How to take TRUQAP + fulvestrant

This dosing schedule was designed to reduce the risk of side effects

TRUQAP is taken 4 days in a row, followed by 3 days not taking TRUQAP, each week.

- Take **TRUQAP** exactly as your doctor tells you
- TRUQAP is taken two times a day about 12 hours apart
- Take your morning and night doses of TRUQAP at the same time each day



The recommended starting dose of TRUQAP is 400 mg two times each day, for a total daily dose of 800 mg. Take TRUQAP exactly as your doctor tells you to, as your doctor may adjust your dose. Do not change your dose or stop taking TRUQAP unless your doctor tells you to.

TRUQAP is taken with fulvestrant

Fulvestrant is an injectable hormone therapy that your doctor will administer on a specific schedule. Each dose of fulvestrant is 500 mg.

In the clinical trial, people received fulvestrant on day 1 and day 15 for cycle 1.

Then, they received fulvestrant on day 1 for the following cycles. Each cycle was 28 days. Your doctor will follow the recommended fulvestrant dosing information in the fulvestrant Prescribing Information.



IMPORTANT SAFETY INFORMATION (continued)

How should I take TRUQAP? (continued)

- Take TRUQAP 2 times each day (about 12 hours apart) in the morning and evening, at about the same times each day
- Take TRUQAP for 4 days in a row, followed by 3 days not taking TRUQAP each week
- Take TRUQAP with or without food

Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.

Convenient packaging to help manage your dosing schedule*

This easy-to-use blister pack that you can take on the go was designed with clear instructions to help you keep track of your medication schedule.

Truqap[®]
capivasertib
160 mg • 200 mg tablets



Travel-friendly packaging



AM/PM color-coded dosing cues




Blank start date on the front for customizable, weekly tracking



Easy-open push tabs

*TRUQAP is available in a blister pack for select doses.



How to store TRUQAP

- Store TRUQAP in original packaging at room temperature between 68 °F to 77 °F (20 °C to 25 °C)
- Throw away (discard) TRUQAP after 45 days if the tablets are not kept in the original packaging
- TRUQAP comes in child-resistant packaging

With TRUQAP, you can choose
the best day of the week to start

Helpful reminders when taking TRUQAP



Swallow whole

Swallow TRUQAP tablets whole with water. Do not chew, crush, or split the tablets



Take with or without food

TRUQAP can be taken with or without food



Don't double up

If you vomit after taking a dose, do not take another dose. Take your next dose at your usual time. Do not take 2 doses at the same time to make up for a missed dose



Do not take damaged tablets

Do not take any tablets that are broken, cracked, or that look damaged



Talk to your doctor

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. TRUQAP may affect the way other medicines work, and other medicines may affect how TRUQAP works

Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.



Avoid grapefruit

You should not drink grapefruit juice or eat grapefruit during treatment with TRUQAP



Accompanying medication

For women who have not reached menopause, your doctor will also prescribe a medicine called a luteinizing hormone-releasing hormone (LHRH) agonist

For men, your healthcare provider may prescribe an LHRH agonist to take with TRUQAP + fulvestrant



Know the medicines you take

Keep a list of all your medications to show your doctor, nurse, or pharmacist when you get a new medicine



Not an actual patient.




If you miss a dose:

If you miss a dose of TRUQAP, you may still take it within 4 hours from the time you usually take it. If it has been more than 4 hours after you usually take your dose, skip that dose. Take the next dose at your usual time. Do not take 2 doses at once to make up for a missed dose.

If you vomit after taking a dose of TRUQAP, do not take an additional dose. Take your next dose at your usual time.

UNDERSTANDING SIDE EFFECTS WITH TRUQAP





Not an actual patient.
The trial showed no significant difference in how long patients lived between treatment groups.



Potential side effects with TRUQAP



TRUQAP may cause side effects, which may sometimes be severe, including



High blood sugar levels

- High blood sugar levels, known as hyperglycemia, are common with TRUQAP and may be severe
- Untreated severe hyperglycemia can lead to a condition called diabetic ketoacidosis, which is a serious condition that requires treatment in a hospital and can lead to death. It is possible to develop diabetic ketoacidosis while taking TRUQAP
- Your doctor will monitor your blood sugar levels before you start and during treatment with TRUQAP
- It is not known if TRUQAP is safe in people with type 1 diabetes or people who use insulin to treat their diabetes
- If you have a history of diabetes, your doctor will monitor your blood levels more often
- Tell your doctor right away if you develop symptoms of high blood sugar, including:
 - excessive thirst
 - dry mouth
 - more frequent urination than usual or a bigger amount of urine than normal
 - blurred vision
 - increased appetite with weight loss
 - stomach area (abdominal) pain
 - unusual tiredness
 - confusion
 - nausea
 - vomiting
 - fruity odor on breath
 - dry or flushed skin
 - difficulty breathing
 - sleepiness



Diarrhea

- Diarrhea is common during treatment and may be severe
- Talk to your doctor if you develop any signs of diarrhea, including loose or watery stool
- Your doctor will tell you to drink more fluids and may prescribe medication to treat diarrhea



Skin reactions

- Skin reactions are common with TRUQAP and may be severe
- Tell your doctor or get medical help right away if you get a new or worsening rash; reddening of the skin; fever; blistering of the lips, eyes, or mouth; blisters on the skin; skin peeling; or dry skin

Your doctor may adjust your treatment if you get certain serious side effects.

Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.




Most common side effects

- nausea
- vomiting
- changes in certain blood tests
- tiredness
- mouth sores



Among patients who experienced these side effects, most side effects appeared within the first 2 weeks of treatment. **See tips for side effect management on pages 16–17.**



Not an actual patient.

Remember to always tell your doctor about any side effects you experience, especially if side effects are severe or if they don't go away or they get worse.


These are not all the possible side effects of TRUQAP. For more information, ask your doctor, nurse, or pharmacist.

You may report side effects to FDA at 1-800-FDA-1088.

Side effect management

There are things you and your healthcare team can do to help

You may experience side effects at any time while taking TRUQAP + fulvestrant. Before treatment even starts, it's important to talk with your doctor or nurse about common side effects, as they may have tips and advice on what to have on hand to make you more comfortable. Tell your doctor about any side effects you may experience. Below are some suggestions of things you may be able to do to manage common side effects.



Tips to Manage Diarrhea

- Call your doctor at the first sign of diarrhea
- Increase fluid intake
- Eat less dairy
- Avoid spicy foods
- Limit caffeine intake
- Taking TRUQAP with food may help reduce the risk of getting diarrhea*



Tips to Manage Nausea

- Eat small meals more often
- Avoid strong smells
- Drink clear, cold beverages
- If you experience nausea, try to avoid foods and drinks that can make you feel worse



Talk to your doctor about any side effects you experience. **There may be things they can do to help you manage certain side effects.**

*Based on an early study in healthy participants.

Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.



Tips to Manage Rashes and Skin Reactions

- At the first sign of skin irritation, call your doctor. Your doctor may recommend over-the-counter creams or prescriptions
- Avoid body washes and lotions with dyes and perfumes
- Switch to dye- and fragrance-free detergent
- Wear cotton or linen fabrics



Tips to Manage Tiredness

- Delegate chores
- Schedule time to relax
- Exercise

Ask your doctor before starting any new physical activity or exercise regimen.

Always tell your doctor about what you are going through, especially if you experience new symptoms or changes.

SUPPORT & RESOURCES FOR TRUQAP





Not an actual patient.
The trial showed no significant difference in how long patients lived between treatment groups.

Support and resources

Helping you access the care you need

If your doctor has prescribed TRUQAP, you may have questions. The AstraZeneca Access 360™ program can help.

We can answer your questions about:



Insurance coverage



Out-of-pocket costs



Patient assistance programs



To learn more about the program, please call

1-844-ASK-A360 (1-844-275-2360)

Monday through Friday, 8 AM – 6 PM ET

or visit www.MyAccess360.com

Explore the Savings

Patients may pay

\$0
co-pay offer

You may be able to **pay as little as \$0 per month with the TRUQAP Patient Savings Program***

*Subject to eligibility. Restrictions apply.

The MyTRUQAP Support Program was created to help you while you're on treatment, with tips and resources, including:



Email updates

Get helpful information and updates about taking TRUQAP and more




TRUQAP resources

Receive helpful resources during treatment



Financial help

Want to learn about savings?
We can help



Not an actual patient.

Scan here to sign up
or visit [TRUQAP.com](https://www.truqap.com)



What is TRUQAP?

TRUQAP is a prescription oral medicine used in combination with the medicine fulvestrant to treat adults who have hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer that is advanced or that has spread to other parts of the body (metastatic), with an abnormal “PIK3CA”, “AKT1”, or “PTEN” gene, and whose disease has progressed on or after endocrine therapy.

Your healthcare provider will test your cancer for an abnormal “PIK3CA”, “AKT1”, or “PTEN” gene to make sure that TRUQAP is right for you.

It is not known if TRUQAP is safe and effective in children.

IMPORTANT SAFETY INFORMATION ABOUT TRUQAP® (capivasertib) tablets

Do not take TRUQAP if you have had a severe allergic reaction to TRUQAP or are allergic to any of the ingredients of TRUQAP.

Before you take TRUQAP, tell your healthcare provider about all your medical conditions, including if you:

- have a history of high levels of sugar in your blood or diabetes
- have a history of rash or other skin disorders
- have liver problems
- have any current infection
- are pregnant or plan to become pregnant. TRUQAP can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider will check to see if you are pregnant before you start treatment with TRUQAP
- You should use effective birth control during treatment with TRUQAP and for 1 month after the last dose. Talk to your healthcare provider about birth control methods that may be right for you during this time
- If you become pregnant or think you are pregnant, tell your healthcare provider right away

Males with female partners who are able to become pregnant should use effective birth control during your treatment with TRUQAP and for 4 months after the last dose. If your female partner becomes pregnant, tell your healthcare provider right away.

- are breastfeeding or plan to breastfeed. Do not breastfeed during treatment with TRUQAP. Talk to your healthcare provider about the best way to feed your baby during treatment with TRUQAP

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. TRUQAP may affect the way other medicines work, and other medicines may affect how TRUQAP works. Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

How should I take TRUQAP?

- Take TRUQAP exactly as your healthcare provider tells you
- Do not change your dose or stop taking TRUQAP unless your healthcare provider tells you
- Take TRUQAP 2 times each day (about 12 hours apart) in the morning and evening, at about the same times each day
- Take TRUQAP for 4 days in a row, followed by 3 days not taking TRUQAP each week

- Take TRUQAP with or without food
- Swallow TRUQAP tablets whole with water. Do not chew, crush, or split the tablets. Do not take any tablets that are broken, cracked, or that look damaged
- If you miss a dose of TRUQAP, you may still take it within 4 hours from the time you usually take it. If it has been more than 4 hours after you usually take your dose, skip that dose. Take the next dose at your usual time. Do not take 2 doses to make up for a missed dose
- If you vomit after taking a dose of TRUQAP, do not take an additional dose. Take your next dose at your usual time
- For women who have not reached menopause or are just starting menopause, your healthcare provider will prescribe a medicine called a luteinizing hormone-releasing hormone (LHRH) agonist to take with TRUQAP and fulvestrant. For men, your healthcare provider may prescribe a LHRH agonist to take with TRUQAP and fulvestrant

What are the possible side effects of TRUQAP?

TRUQAP may cause severe side effects, including:

- **High blood sugar levels (hyperglycemia)** – Hyperglycemia is common with TRUQAP and may be severe. Untreated severe hyperglycemia can lead to a condition called diabetic ketoacidosis that can happen in people treated with TRUQAP. Diabetic ketoacidosis is a serious condition that requires treatment in a hospital and that can lead to death. Your healthcare provider will monitor your blood sugar levels before you start and during treatment with TRUQAP. It is not known if TRUQAP is safe in people with type 1 diabetes or people who use insulin to treat their diabetes. Your healthcare provider will monitor your blood sugar levels more often if you have a history of diabetes. Tell your healthcare provider right away if you develop symptoms of hyperglycemia, including excessive thirst, dry mouth, more frequent urination than usual or a bigger amount of urine than normal, blurred vision, increased appetite with weight loss, stomach area (abdominal) pain, unusual tiredness, confusion, nausea, vomiting, fruity odor on breath, dry or flushed skin, difficulty breathing, sleepiness
- **Diarrhea** – Diarrhea is common during treatment with TRUQAP and may be severe. Severe diarrhea can lead to the loss of too much body water (dehydration). Tell your healthcare provider if you develop any signs of diarrhea, including loose or watery stool. Your healthcare provider will tell you to drink more fluids or take medicines to treat diarrhea
- **Skin reactions** – Skin reactions are common with TRUQAP and can be severe. Tell your healthcare provider or get medical help right away if you get a new or worsening rash, reddening of the skin, fever, blistering of the lips, eyes or mouth, blisters on the skin, skin peeling, or dry skin

Your healthcare provider may tell you to decrease your dose, temporarily stop your treatment, or completely stop your treatment with TRUQAP if you get certain serious side effects.

The most common side effects of TRUQAP include nausea, tiredness, vomiting, mouth sores and changes in certain blood tests.

These are not all the possible side effects of TRUQAP. Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

What should I avoid while taking TRUQAP?

- You should not drink grapefruit juice or eat grapefruit during treatment with TRUQAP

Please see full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.

You may [report side effects related to AstraZeneca products](#) .



The trial showed no significant difference in how long patients lived between treatment groups.

UNFOLDING MORE OF YOUR STORY MAY BE POSSIBLE*

MORE THAN **2X** longer without cancer growing or spreading*

***TRUQAP + fulvestrant** delivered an average of 7.3 months vs 3.1 months with fulvestrant alone—a difference of 4.2 months. Results may vary.

See helpful tips to manage potential side effects inside

Scan here to learn more about TRUQAP



IMPORTANT SAFETY INFORMATION ABOUT TRUQAP® (capivasertib) tablets

Do not take TRUQAP if you have had a severe allergic reaction to TRUQAP or are allergic to any of the ingredients of TRUQAP.

Before you take TRUQAP, tell your healthcare provider about all your medical conditions, including if you:

- have a history of high levels of sugar in your blood or diabetes
- have a history of rash or other skin disorders

Please see additional Important Safety Information throughout, and full Prescribing Information, including Patient Information for TRUQAP.

AstraZeneca 

TRUQAP is a registered trademark and AstraZeneca Access 360 is a trademark of the AstraZeneca group of companies.

©2026 AstraZeneca. All rights reserved. US-112077 Last Updated 4/26