

TRUQAP + abiraterone with prednisone  
is for certain people with metastatic\*  
prostate cancer that is PTEN-deficient

# MAN WITH A PLAN AND TRUQAP

The **FIRST AND ONLY** targeted  
combination therapy of its kind

\*Metastatic=cancer that has spread to other  
parts of the body.

Actor portrayals.

## IMPORTANT SAFETY INFORMATION ABOUT TRUQAP® (capivasertib) tablets

**Do not take TRUQAP** if you have had a severe allergic reaction to TRUQAP or are allergic to any of the ingredients of TRUQAP.

**Before you take TRUQAP, tell your healthcare provider about all your medical conditions, including if you:**

- have a history of high levels of sugar in your blood or diabetes
- have a history of rash or other skin disorders
- have liver problems
- have any current infection

**Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.**

 **Truqap®**  
capivasertib  
160 mg • 200 mg tablets

## Glossary

Throughout this brochure, you will see abbreviations and terms associated with metastatic prostate cancer (mPC).

Here's what they mean:

**Abiraterone**=an oral hormone therapy drug taken with prednisone (steroid)

**ADT**=a type of hormone therapy used for prostate cancer (stands for “androgen deprivation therapy”)

**Biomarkers**=signs in your body that doctors can find to learn more about your health

**GnRH**=gonadotropin-releasing hormone

**IHC**=a test done on the tissue sample taken through a biopsy to detect PTEN deficiency in tumors (stands for “immunohistochemistry”)

**Metastatic**=cancer that has spread to other parts of the body

**mHSPC**=metastatic hormone-sensitive prostate cancer

**mPC**=metastatic prostate cancer

**Prednisone**=a steroid medication

**PTEN**=a protein that helps block cancer signals (stands for “phosphatase and tensin homolog”)

**PTEN deficiency**=a lack of PTEN, a protein that blocks cancer signals. Your doctor or healthcare team may also refer to it as PTEN loss

**rPFS**=a measure of how long a treatment keeps cancer from growing or spreading (stands for “radiographic progression-free survival”)



## Table of contents

|                                                     |               |
|-----------------------------------------------------|---------------|
| <b>Glossary .....</b>                               | <b>pg. 1</b>  |
| <b>mPC and PTEN deficiency .....</b>                | <b>pg. 3</b>  |
| <b>Start your plan with biomarker testing .....</b> | <b>pg. 4</b>  |
| <b>How TRUQAP works .....</b>                       | <b>pg. 6</b>  |
| <b>Results with TRUQAP .....</b>                    | <b>pg. 7</b>  |
| <b>How TRUQAP was studied .....</b>                 | <b>pg. 8</b>  |
| <b>How to take TRUQAP .....</b>                     | <b>pg. 9</b>  |
| <b>Helpful reminders when taking TRUQAP .....</b>   | <b>pg. 11</b> |
| <b>Potential side effects with TRUQAP .....</b>     | <b>pg. 13</b> |
| <b>Side effect management .....</b>                 | <b>pg. 15</b> |
| <b>Support and resources .....</b>                  | <b>pg. 17</b> |
| <b>My plan .....</b>                                | <b>pg. 19</b> |
| <b>Important Safety Information .....</b>           | <b>pg. 21</b> |

## Metastatic\* prostate cancer and PTEN deficiency

TRUQAP + abiraterone with prednisone is for certain people with metastatic prostate cancer that is PTEN-deficient

If you or your loved one has been diagnosed with metastatic prostate cancer, that means the disease has spread to other parts of the body.

- For certain people, their metastatic\* prostate cancer responds to hormone therapy, a treatment that reduces hormone levels to help manage their cancer. This may be referred to as metastatic hormone-sensitive prostate cancer, or mHSPC
- However, when tumors have PTEN deficiency, there is an additional path for the tissue to grow despite hormone therapy
- Because of this, combining hormone therapy with another treatment that specifically targets PTEN deficiency may be right for certain people with metastatic prostate cancer that is PTEN-deficient



Actor portrayals.

### What is PTEN deficiency?

PTEN deficiency refers to a lack of PTEN, a protein that blocks cancer signals. When PTEN is lacking, tumors grow more quickly and aggressively. Your doctor or healthcare team may also refer to it as PTEN loss.

\*Metastatic=cancer that has spread to other parts of the body.



**A biomarker test can determine if you have PTEN deficiency**



## Start your plan with biomarker testing

Biomarkers are signs in your body that doctors can find to learn more about your health. PTEN deficiency is a biomarker that can change how tumors grow in metastatic\* prostate cancer.

- PTEN deficiency is a biomarker that may cause tumors to grow more quickly
- Around 25% of people with metastatic\* prostate cancer that is responding to hormone therapy have tumors with PTEN deficiency
- To test for PTEN deficiency (with IHC), your doctor uses a small tissue sample of the tumor that is taken through a biopsy
- Testing for biomarkers like PTEN deficiency can help your doctor figure out if a targeted treatment may be right for you

**TRUQAP + abiraterone,<sup>†</sup> a targeted treatment option, is the first combination therapy of its kind for certain people with metastatic prostate cancer that is PTEN-deficient.**

<sup>†</sup>Abiraterone is taken with prednisone.



Actor portrayals.



**The TRUQAP plan starts with talking to your doctor about testing for PTEN deficiency**

### IMPORTANT SAFETY INFORMATION (continued)

**Males** with female partners who are able to become pregnant should use effective birth control during your treatment with TRUQAP and for 4 months after the last dose. If your female partner becomes pregnant, tell your healthcare provider right away.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. TRUQAP may affect the way other medicines work, and other medicines may affect how TRUQAP works. Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

**Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.**







## How TRUQAP works

with abiraterone + prednisone

### The first targeted combination therapy of its kind

TRUQAP is taken in combination with a hormone therapy called abiraterone.\* These medicines work together to block sources of cancer growth.

In metastatic<sup>†</sup> prostate cancer with PTEN deficiency, there are two different sources in the body that cause cancer cells to grow and spread. TRUQAP + abiraterone\* work together by blocking both sources of tumor growth at the same time.

- TRUQAP blocks cancer signals in one of those sources from telling cancer cells to grow and spread
- Abiraterone\* lowers testosterone, a hormone that fuels tumor growth

This combination targets two important sources of cancer growth.

\*Abiraterone is taken with prednisone.

<sup>†</sup>Metastatic=cancer that has spread to other parts of the body.

### IMPORTANT SAFETY INFORMATION (continued)

#### How should I take TRUQAP?

- Take TRUQAP exactly as your healthcare provider tells you
- Do not change your dose or stop taking TRUQAP unless your healthcare provider tells you
- Take TRUQAP 2 times each day (about 12 hours apart) in the morning and evening, at about the same times each day
- Take TRUQAP for 4 days in a row, followed by 3 days not taking TRUQAP each week
- Take TRUQAP with or without food
- Swallow TRUQAP tablets whole with water. Do not chew, crush, or split the tablets. Do not take any tablets that are broken, cracked, or that look damaged

Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.



## Results with TRUQAP in combination with abiraterone + prednisone

TRUQAP in combination\* helped people get more time without their cancer growing or spreading vs abiraterone + prednisone.

TRUQAP in combination\* delivered a median of 33.2 months without their cancer growing or spreading vs 25.7 months with abiraterone + prednisone—a difference of 7.5 months. Results may vary.



Radiographic progression-free survival (rPFS) is a measure of how long a treatment keeps cancer from growing or spreading. Median rPFS is the time point at which half the patients have experienced disease progression. Median is the middle number in a group of numbers listed from lowest to highest.

\*Combination therapy: TRUQAP + abiraterone + prednisone.

†Abiraterone was taken with prednisone.

### IMPORTANT SAFETY INFORMATION (continued)

#### How should I take TRUQAP? (continued)

- If you miss a dose of TRUQAP, you may still take it within 4 hours from the time you usually take it. If it has been more than 4 hours after you usually take your dose, skip that dose. Take the next dose at your usual time. Do not take 2 doses to make up for a missed dose
- If you vomit after taking a dose of TRUQAP, do not take an additional dose. Take your next dose at your usual time
- For men who have not had surgery to remove both testicles (bilateral orchiectomy), your healthcare provider will prescribe a gonadotropin-releasing hormone (GnRH) to take with TRUQAP along with abiraterone and prednisone

Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.

## How TRUQAP was studied

TRUQAP was studied in a global clinical trial that included 1,012 people with metastatic\* prostate cancer and PTEN deficiency.

### The study group included:

**1,012**  
People

**507**

TRUQAP +  
abiraterone +  
prednisone†

**505**

Abiraterone  
+ prednisone†

- For certain people with metastatic\* prostate cancer that is PTEN-deficient
- Tumors with PTEN deficiency (confirmed with an IHC test)
- Median age of 68

### Two of the study's main goals were:



See how well the combination therapy‡ could keep cancer from growing or spreading vs abiraterone + prednisone



Determine the safety of TRUQAP in combination‡

\*Metastatic=cancer that has spread to other parts of the body.

†All patients were also taking ADT.

‡Combination therapy: TRUQAP + abiraterone + prednisone.



**Talk to your doctor about getting tested for PTEN deficiency**



# How to take TRUQAP + abiraterone + prednisone

TRUQAP



**4 DAYS ON / 3 DAYS OFF**

Two tablets (200 mg each), twice a day, totaling 4 tablets each day by mouth (800 mg daily dose), 4 days a week.

Take TRUQAP exactly as your doctor tells you to, as your doctor may adjust your dose. Do not change your dose or stop taking TRUQAP unless your doctor tells you to.

TRUQAP is taken in combination with other medicines. Each has its own dosing schedule.



**ABIRATERONE (ABI)**

Two tablets (500 mg each), once a day (1,000 mg daily dose), regardless of whether it is a day you are taking TRUQAP or not.



**PREDNISONE (PRED)**

Your doctor will also prescribe a steroid, prednisone. 5 mg taken orally, once daily, or according to how your doctor has prescribed.



Your doctor will also prescribe ADT. Take it according to how your doctor has prescribed.

If you have not had surgery to remove both testicles (bilateral orchiectomy), your healthcare provider will prescribe a GnRH to take with TRUQAP. This will be in addition to abiraterone and prednisone.

| Treatment schedule | DAY 1                           | DAY 2                           | DAY 3                           | DAY 4                           | DAY 5                           | DAY 6                           | DAY 7                           |
|--------------------|---------------------------------|---------------------------------|---------------------------------|---------------------------------|---------------------------------|---------------------------------|---------------------------------|
| AM                 | TRUQAP<br>+<br>ABI<br>+<br>PRED | TRUQAP<br>+<br>ABI<br>+<br>PRED | TRUQAP<br>+<br>ABI<br>+<br>PRED | TRUQAP<br>+<br>ABI<br>+<br>PRED | TRUQAP<br>+<br>ABI<br>+<br>PRED | TRUQAP<br>+<br>ABI<br>+<br>PRED | TRUQAP<br>+<br>ABI<br>+<br>PRED |
| PM                 | TRUQAP                          | TRUQAP                          | TRUQAP                          | TRUQAP                          | TRUQAP                          | TRUQAP                          | TRUQAP                          |

## Blister packs designed to help you manage your dosing schedule

This blister pack that you can take on the go was designed with clear instructions to help you keep track of your medication schedule.



Blank start date on the front for customizable, weekly tracking



AM/PM-labeled and color-coded dosing cues



Travel-friendly packaging

## How should I store TRUQAP?

- Store TRUQAP in original packaging at room temperature between 68 °F to 77 °F (20 °C to 25 °C)
- Throw away (discard) TRUQAP after 45 days if the tablets are not kept in the original packaging
- TRUQAP comes in child-resistant packaging

## Did you miss a dose?

If you miss a dose of TRUQAP, you may still take it within 4 hours from the time you usually take it. If it has been more than 4 hours after you usually take your dose, skip that dose. Take the next dose at your usual time. Do not take 2 doses to make up for a missed dose.



With TRUQAP, you can choose the best day of the week to start



## Helpful reminders when taking TRUQAP



### Do not take damaged tablets

Do not take any tablets that are broken, cracked, or that look damaged



### Avoid grapefruit

You should not drink grapefruit juice or eat grapefruit during treatment with TRUQAP



### Swallow whole

Swallow TRUQAP tablets whole with water. Do not chew, crush, or split the tablets



### Take with or without food

TRUQAP can be taken with or without food




### Don't double up

If you vomit after taking a dose, do not take another dose. Take your next dose at your usual time



### Talk to your doctor

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. TRUQAP may affect the way other medicines work, and other medicines may affect how TRUQAP works



### Know the medicines you take


Keep a list of them to show your doctor, nurse, or pharmacist when you get a new medicine





# Potential side effects with TRUQAP

TRUQAP may cause side effects, which may sometimes be severe, including:



## High blood sugar levels

- High blood sugar levels, known as hyperglycemia, are common with TRUQAP and can be severe
- Hyperglycemia has happened in people with and without a history of diabetes
- Untreated severe hyperglycemia can lead to a condition called diabetic ketoacidosis, which is a serious condition that requires treatment in a hospital and can lead to death. It is possible to develop diabetic ketoacidosis while taking TRUQAP
- **Your doctor will monitor your blood sugar levels before you start and during treatment with TRUQAP**
- It is not known if TRUQAP is safe in people with type 1 diabetes or people who use insulin to treat their diabetes
- If you have a history of diabetes, your doctor will monitor your blood sugar levels more often
- If you have a fever, chills, infection, serious injury, or new illness, tell your healthcare provider before taking your next dose
- Tell your doctor right away if you develop symptoms of high blood sugar, including:
  - excessive thirst
  - dry mouth
  - more frequent urination than usual or a bigger amount of urine than normal
  - blurred vision
  - increased appetite with weight loss
  - stomach area (abdominal) pain
  - unusual tiredness
  - confusion
  - nausea
  - vomiting
  - fruity odor on breath
  - dry or flushed skin
  - difficulty breathing
  - sleepiness

Your doctor may adjust your treatment if you experience certain serious side effects.

Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.



## Diarrhea

- Diarrhea is common during treatment and can be severe
- Talk to your doctor if you develop any signs of diarrhea, including loose or watery stool
- Your doctor will tell you to drink more fluids and may prescribe medication to treat diarrhea



## Skin reactions

- Skin reactions are common with TRUQAP and can be severe
- Tell your doctor or get medical help right away if you experience a new or worsening rash; reddening of the skin; a fever; blistering of the lips, eyes, or mouth; blisters on the skin; skin peeling; or dry skin

## Most common side effects:

- Skin reactions
- Diarrhea
- Tiredness
- Changes in certain blood tests

Be sure to tell your doctor if you experience any side effects, even if they appear minor. It's important to address them before they get worse. These are not all the possible side effects of TRUQAP. For more information, ask your doctor, nurse, or pharmacist. You may report side effects to FDA at 1-800-FDA-1088.



Among people who experienced rash and diarrhea, these symptoms usually appeared within the first 2 weeks of treatment. **See tips for side effect management on pages 15 and 16**



## Side effect management

There are things you and your healthcare team can do to help

You may experience side effects at any time while taking TRUQAP + abiraterone with prednisone. Before treatment even starts, it's important to talk about common side effects with your doctor or nurse, as they may have tips for making yourself more comfortable. Tell your doctor about any side effects you may experience. Below are some suggestions that may help you manage common side effects.



### Tips to help manage diarrhea

- Call your doctor at the first sign of diarrhea
- Increase fluid intake
- Take medicines to treat diarrhea
- Eat less dairy
- Avoid spicy foods
- Limit caffeine intake
- Taking TRUQAP with food may help reduce the risk of getting diarrhea\*



### Tips to help manage rashes and skin reactions

- At the first sign of skin irritation, call your doctor. Your doctor may recommend over-the-counter creams or prescriptions
- Avoid body washes and lotions with dyes and perfumes
- Switch to dye- and fragrance-free detergent
- Wear cotton or linen fabrics



**If you experience any side effects, talk to your doctor right away**

\*Based on an early study in healthy participants.




### Tips to help manage nausea

- Eat small meals more often
- Avoid strong smells
- Drink clear, cold beverages
- If you experience nausea, try to avoid foods and drinks that can make you feel worse



### Tips to help manage tiredness

- Delegate chores
- Schedule time to relax
- Exercise
- Ask your doctor before starting any new physical activity or exercise regimen



Actor portrayal.



Scan here for a helpful  
side effect tracker.

## Support and resources

### Helping you access the care you need

If your doctor has prescribed TRUQAP, you may have questions. The AstraZeneca Access 360™ program can help.

### We can answer your questions about:



Insurance coverage



Out-of-pocket costs



Patient assistance programs



To learn more about the program, please call  
1-844-ASK-A360 (1-844-275-2360)  
Monday through Friday, 8 AM – 6 PM ET  
or visit [www.MyAccess360.com](http://www.MyAccess360.com)

#### Explore the Savings

Patients may pay

**\$0\***  
co-pay offer

You may be able to **pay as little as \$0 per month with the TRUQAP Patient Savings Program\***

\*Subject to eligibility. Restrictions apply.



The **MyTRUQAP Support Program** was created to help you while you're on treatment, with tips and resources, including:



### Email updates

Get helpful information and updates about taking TRUQAP and more



### TRUQAP resources

Receive helpful resources during treatment



### Financial help

Want to learn about savings?  
We can help



Actor portrayals.



Scan to visit [TRUQAP.com/patient-support-program](https://truqap.com/patient-support-program) and sign up for more Patient and Care Partner support.

## My plan

Keep track of your treatment notes on this page.

[illegible]







## What is TRUQAP?

TRUQAP is a prescription oral medicine used in combination with the medicines abiraterone and prednisone to treat adults who have PTEN-deficient prostate cancer that has spread to other parts of the body (metastatic).

Your healthcare provider will test your prostate cancer to see if it has PTEN deficiency to make sure that TRUQAP is right for you.

It is not known if TRUQAP is safe and effective in children.

## IMPORTANT SAFETY INFORMATION ABOUT TRUQAP® (capivasertib) tablets

**Do not take TRUQAP if you** have had a severe allergic reaction to TRUQAP or are allergic to any of the ingredients of TRUQAP.

**Before you take TRUQAP, tell your healthcare provider about all your medical conditions, including if you:**

- have a history of high levels of sugar in your blood or diabetes
- have a history of rash or other skin disorders
- have liver problems
- have any current infection

**Males** with female partners who are able to become pregnant should use effective birth control during your treatment with TRUQAP and for 4 months after the last dose. If your female partner becomes pregnant, tell your healthcare provider right away.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. TRUQAP may affect the way other medicines work, and other medicines may affect how TRUQAP works. Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

## How should I take TRUQAP?

- Take TRUQAP exactly as your healthcare provider tells you
- Do not change your dose or stop taking TRUQAP unless your healthcare provider tells you
- Take TRUQAP 2 times each day (about 12 hours apart) in the morning and evening, at about the same times each day
- Take TRUQAP for 4 days in a row, followed by 3 days not taking TRUQAP each week
- Take TRUQAP with or without food
- Swallow TRUQAP tablets whole with water. Do not chew, crush, or split the tablets. Do not take any tablets that are broken, cracked, or that look damaged
- If you miss a dose of TRUQAP, you may still take it within 4 hours from the time you usually take it. If it has been more than 4 hours after you usually take your dose, skip that dose. Take the next dose at your usual time. Do not take 2 doses to make up for a missed dose
- If you vomit after taking a dose of TRUQAP, do not take an additional dose. Take your next dose at your usual time
- For men who have not had surgery to remove both testicles (bilateral orchiectomy), your healthcare provider will prescribe a gonadotropin-releasing hormone (GnRH) to take with TRUQAP along with abiraterone and prednisone

### What are the possible side effects of TRUQAP?

#### TRUQAP may cause severe side effects, including:

- **High blood sugar levels (hyperglycemia)** – Hyperglycemia is common with TRUQAP and can be severe. Hyperglycemia has happened in people with and without a history of diabetes. Untreated severe hyperglycemia can lead to a condition called diabetic ketoacidosis that can happen in people treated with TRUQAP. Diabetic ketoacidosis is a serious condition that requires treatment in a hospital and that can lead to death. Your healthcare provider will monitor your blood sugar levels before you start and during treatment with TRUQAP. It is not known if TRUQAP is safe in people with type 1 diabetes or people who use insulin to treat their diabetes. Your healthcare provider will monitor your blood sugar levels more often if you have a history of diabetes. If you have a fever, chills, infection, serious injury, or a new illness, tell your healthcare provider before taking your next dose. Tell your healthcare provider right away if you develop symptoms of hyperglycemia, including excessive thirst, dry mouth, more frequent urination than usual or a bigger amount of urine than normal, blurred vision, increased appetite with weight loss, stomach area (abdominal) pain, unusual tiredness, confusion, nausea, vomiting, fruity odor on breath, dry or flushed skin, difficulty breathing, sleepiness
- **Diarrhea** – Diarrhea can be severe during treatment with TRUQAP. Severe diarrhea can lead to the loss of too much body water (dehydration). Tell your healthcare provider if you develop any signs of diarrhea, including loose or watery stool. Your healthcare provider will tell you to drink more fluids or take medicines to treat diarrhea
- **Skin reactions** – Skin reactions can be severe during treatment with TRUQAP. Tell your healthcare provider or get medical help right away if you get a new or worsening rash, reddening of the skin, fever, blistering of the lips, eyes or mouth, blisters on the skin, skin peeling, or dry skin

Your healthcare provider may tell you to decrease your dose, temporarily stop your treatment, or completely stop your treatment with TRUQAP if you get certain serious side effects.

**The most common side effects of TRUQAP when used in combination with abiraterone and prednisone include** skin reactions, diarrhea, tiredness, and changes in certain blood tests.

These are not all the possible side effects of TRUQAP. Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

### What should I avoid while taking TRUQAP?

- You should not drink grapefruit juice or eat grapefruit during treatment with TRUQAP

**Please see full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.**

You may [report side effects related to AstraZeneca products](#) .

## Here's the plan



Follow the dosing schedule



Let your doctor know about any side effects



Get financial support with the AstraZeneca Access 360™ program



Learn more at  
[TRUQAP.com/prostate-cancer](https://truqap.com/prostate-cancer)

Actor portrayal.

### IMPORTANT SAFETY INFORMATION ABOUT TRUQAP® (capivasertib) tablets

**Do not take TRUQAP if you** have had a severe allergic reaction to TRUQAP or are allergic to any of the ingredients of TRUQAP.

**Before you take TRUQAP, tell your healthcare provider about all your medical conditions, including if you:**

- have a history of high levels of sugar in your blood or diabetes
- have a history of rash or other skin disorders
- have liver problems
- have any current infection

**Please see additional Important Safety Information throughout, and full Prescribing Information, including Patient Information for TRUQAP.**



TRUQAP is a registered trademark and AstraZeneca Access 360 is a trademark of the AstraZeneca group of companies.  
©2026 AstraZeneca. All rights reserved. US-107454 Last Updated 6/26