

Recognizing Side Effects Early

*Metastatic=cancer that has spread to other parts of the body.

Did you check in with your body today to identify any potential side effects?

Side effects may occur while taking TRUQAP + abiraterone with prednisone, including those that can be severe.

What to look out for and what to do if you experience symptoms



High blood sugar levels

What to look out for:

If you have a fever, chills, infection, serious injury, or a new illness, tell your healthcare provider before taking your next dose.

Symptoms of high blood sugar include:

- excessive thirst
- dry mouth
- more frequent urination than usual or a bigger amount of urine than normal
- blurred vision
- increased appetite with weight loss
- stomach area (abdominal) pain
- unusual tiredness
- confusion
- nausea or vomiting
- fruity odor on breath
- dry or flushed skin
- difficulty breathing
- sleepiness

What to do:

- Your doctor will monitor your blood sugar levels before you start and during treatment with TRUQAP
- Your doctor may recommend lifestyle changes like eating a lower-carbohydrate, high-fiber diet and staying active regularly

Untreated severe hyperglycemia can lead to a condition called diabetic ketoacidosis, which is a serious condition that requires treatment in a hospital and can lead to death.



Skin reactions

What to look out for:

- a rash
- a rash that keeps getting worse
- reddening of the skin
- fever
- blistering of the lips, eyes, or mouth
- blisters on the skin
- dry or peeling skin

What to do:

- Avoid body washes and lotions with dyes and perfumes
- Switch to dye- and fragrance-free detergent
- Wear cotton or linen fabrics
- Your doctor may recommend over-the-counter creams or prescriptions. Keep them handy, and take as advised if skin reactions occur

Watch out for signs of an allergic reaction, such as swelling, chest pain, or difficulty breathing, as well as symptoms consistent with an allergic reaction, including fever, chills, and rash.



Diarrhea

What to look out for:

- loose or watery stool
- increase in bowel movements

What to do:

- Increase fluid intake
- Limit dairy products, avoid spicy food, and limit caffeine intake
- Taking TRUQAP with food may help reduce the risk of getting diarrhea[†]
- Use your Side Effect Tracker to log bowel movements, stool changes, and any antidiarrheal use. Your doctor can use this information to help manage your diarrhea
- Your doctor may suggest some over-the-counter treatments that you could have on hand to help with diarrhea. Keep them handy and take as advised

[†]Based on an early study in healthy participants.

As soon as you notice any symptoms, call your healthcare team right away. They can advise you regarding these symptoms. Then record them in your Side Effect Tracker.



These are not all the possible side effects with TRUQAP + abiraterone with prednisone. Your doctor may adjust your treatment or prescribe medication to help manage your side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Please see full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.

Recognizing Side Effects Early (cont'd)

My healthcare team

Name: _____

Phone: _____

Address: _____

Website: _____

Name: _____

Phone: _____

Address: _____

Website: _____

Name: _____

Phone: _____

Address: _____

Website: _____

Name: _____

Phone: _____

Address: _____

Website: _____

My notes

IMPORTANT PRODUCT INFORMATION

TRUQAP is a prescription oral medicine used in combination with the medicines abiraterone and prednisone to treat adults who have PTEN-deficient prostate cancer that has spread to other parts of the body (metastatic). Do not take TRUQAP if you have had a severe allergic reaction to TRUQAP or are allergic to any of its ingredients. Males with female partners who are able to become pregnant should use effective birth control during your treatment with TRUQAP and for 4 months after the last dose. If your female partner becomes pregnant, tell your healthcare provider right away. TRUQAP may cause severe side effects, including high blood sugar levels (hyperglycemia); diarrhea; or skin reactions. Untreated severe hyperglycemia can lead to a condition called diabetic ketoacidosis that requires treatment in a hospital and can lead to death. The most common side effects include skin reactions, diarrhea, tiredness, and changes in certain blood tests.

Please see full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.