

 Daily Side Effect Tracker
An everyday tool to keep track of side effects

*Metastatic=cancer that has spread to other parts of the body.

This **Daily Side Effect Tracker** is to record and report any side effects experienced while taking TRUQAP + abiraterone with prednisone. The Daily Side Effect Tracker is a convenient complement to the larger Side Effect Tracker. Bring it to your next appointment to inform the healthcare team about any side effects experienced to help keep treatment on track.

Keep track of symptoms of high blood sugar

Tell your doctor right away if you develop symptoms of high blood sugar, known as hyperglycemia, including:

 excessive thirst, dry mouth, more frequent urination than usual or a bigger amount of urine than normal, blurred vision, increased appetite with weight loss, stomach area (abdominal) pain, unusual tiredness, confusion, nausea, vomiting, fruity odor on breath, dry or flushed skin, difficulty breathing, sleepiness If you have a fever, chills, infection, serious injury, or a new illness, tell your healthcare provider before taking your next dose	Date symptom(s) occurred

Keep track of occurrences of diarrhea

 Among patients in the clinical trial who experienced diarrhea, symptoms usually appeared within the first 2 weeks of treatment: - Diarrhea is common during treatment and can be severe - Severe diarrhea can lead to the loss of too much body water (dehydration) - Talk to your doctor if you develop any signs of diarrhea, including loose or watery stool. Your doctor may suggest increasing fluid intake - Ask your doctor about over-the-counter treatments that you could have on hand that may help if you have diarrhea	Date	Number of bowel movements today	Are you experiencing diarrhea?
			☐
			☐
			☐
			☐
			☐
			☐
			☐

IMPORTANT PRODUCT INFORMATION

TRUQAP is a prescription oral medicine used in combination with the medicines abiraterone and prednisone to treat adults who have PTEN-deficient prostate cancer that has spread to other parts of the body (metastatic). Do not take TRUQAP if you have had a severe allergic reaction to TRUQAP or are allergic to any of its ingredients. Males with female partners who are able to become pregnant should use effective birth control during your treatment with TRUQAP and for 4 months after the last dose. If your female partner becomes pregnant, tell your healthcare provider right away. (continued on other side)

Please see full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.

TRUQAP + abiraterone with prednisone is for certain people with metastatic* prostate cancer that is PTEN-deficient



Daily Side Effect Tracker

An everyday tool to keep track of side effects



*Metastatic=cancer that has spread to other parts of the body.

Keep track of any skin reactions, which are common with TRUQAP and can be severe

Tell your doctor or get medical help right away if you develop any of these symptoms:



a new or worsening rash; reddening of the skin; fever; blistering of the lips, eyes, or mouth; blisters on the skin; skin peeling; dry skin

Date symptom(s) occurred

Keep track of other possible side effects

Tell your doctor if you experience any of these common side effects of TRUQAP:

- tiredness
- changes in certain blood tests

Keep track of other possible side effects. **Watch out for signs of an allergic reaction, such as swelling, chest pain, or difficulty breathing.**

Remember to always tell the doctor right away about any side effects experienced. The doctor may adjust treatment or prescribe medication to help manage side effects. With certain serious side effects the doctor may decrease your dose, temporarily stop your treatment, or completely stop your treatment with TRUQAP. These are not all the possible side effects with TRUQAP + abiraterone with prednisone.

You may report side effects to the FDA at 1-800-FDA-1088.

Description of symptom(s)

Date symptom(s) occurred

IMPORTANT PRODUCT INFORMATION (continued)

TRUQAP may cause severe side effects, including high blood sugar levels (hyperglycemia); diarrhea; or skin reactions. Untreated severe hyperglycemia can lead to a condition called diabetic ketoacidosis that requires treatment in a hospital and can lead to death. The most common side effects include skin reactions, diarrhea, tiredness, and changes in certain blood tests.

Please see full [Prescribing Information](#), including [Patient Information](#) for TRUQAP.